

SEXUALLY TRANSMITTED INFECTION FOLLOW-UP GONORRHEA

Gonorrhea is a reportable infection under Ontario Regulation 559/91, as amended, under the Health Protection and Promotion Act. This information is required for case investigation/management and local surveillance.

INSTRUCTIONS: CONFIRM CLIENT DEMOGRAPHICS and return **COMPLETE FORM** via confidential fax to the EOHU at 613-933-7417

Diagnosing Clinician: _____ Date of Positive Test: _____
(YYYY/MM/DD)

Form Completed By (print): _____ Fax Return Form to: _____

CLIENT DEMOGRAPHICS - CONFIRMED:

Name: _____ DOB: _____ Sex: _____
(YYYY/MM/DD)

Address: _____

Email: _____ Phone: _____

TREATMENT (as per Public Health Ontario Gonorrhea Testing and Treatment Guide, 2nd Edition)

FIRST-LINE TREATMENT

(Test of cure recommended)

Ceftriaxone 500mg IM Single Dose

Date Prescribed (YYYY/MM/DD): _____

SECOND-LINE TREATMENT (Perform test of cure)

Alternative treatment for anogenital infections

Cefixime 800mg PO + Doxycycline 100mg PO BID for 7 Days

• This regimen is recommended if there is macrolide resistance or contraindication to macrolide use.

• Doxycycline is contraindicated in pregnant and lactating individuals

Date Prescribed (YYYY/MM/DD): _____

Alternative treatment for pharyngeal infections

Cefixime 800mg PO + Azithromycin 1g PO Single Dose

Date Prescribed (YYYY/MM/DD): _____

COMPLETED CLIENT COUNSELLING

- Notified of positive result
- Advised to abstain from sexual contact for 7 days after treatment
- Advised to abstain from sexual contact with untreated sexual contact(s)
- Informed that retreatment required if vomiting occurs within 1 hour of taking Azithromycin
- Recommended test of cure*

***A test of cure (TOC) is recommended for all positive NG sites in all cases.**

This is particularly important when regimens other than ceftriaxone 500 mg IM are used. Refer to the Test of cure section in the Gonorrhea Guide for more information on the timing for TOC.

- If test of cure is done 3-21 days after treatment completion, culture is recommended regardless of the patient's symptom status
- If test of cure is done after 21 days from treatment completion and the patient is
 - Asymptomatic-NAAT are recommended
 - Symptomatic-Both culture and NAAT are recommended

CONTACT TRACING (from past 3 months or last know if greater than 3 months)

- Client advised to notify sexual contact(s) of need for STI testing and treatment; **AND**
- Client advised EOHU will contact them to collect sexual contact(s) information and complete contact tracing

RISK FACTORS

- | | | |
|--|--|--|
| <input type="checkbox"/> Condom breakage | <input type="checkbox"/> No condom used | <input type="checkbox"/> Same sex partner(s) |
| <input type="checkbox"/> Diagnosed at time of IUD insertion | <input type="checkbox"/> Opposite sex partner(s) | <input type="checkbox"/> Sex trade worker |
| <input type="checkbox"/> Diagnosed at time of therapeutic abortion | <input type="checkbox"/> Parent of a neonate | <input type="checkbox"/> Sexually assaulted |
| <input type="checkbox"/> More than 1 sexual contact in last 6 months | <input type="checkbox"/> Pregnant (Number of weeks: _____) | <input type="checkbox"/> Other: _____ |
| <input type="checkbox"/> New sexual contact in past 2 months | <input type="checkbox"/> On PrEP | |

REASON FOR TESTING

- Contact tracing (sexual contact tested positive)
- Routine
- Symptoms: _____

NOTES

THE EOHU WILL CONTACT ALL CLIENTS INFECTED WITH GONORRHEA.



EOHU.ca • 613-933-1375 • 1-800-267-7120



If you require this information in an alternate format, please call 1-800-267-7120 and press 0.