SEXUALLY TRANSMITTED INFECTION FOLLOW-UP
GONORRHEA

Gonorrhea is a reportable infection under Ontario Regulation 559/91, as amended, under the Health Protection and Promotion Act.
This information is required for case investigation/management and local surveillance.

INSTRUCTIONS: CONFIRM CLIENT DEMOGRAPHICS and return **COMPLETE FORM** via confidential fax to the EOHU at 613-933-7417

Diagnosing Clinician: _____ Date of Positive Test: _____ (YYYY/MM/DD)

Form Completed By (print): _____ Fax Return Form to: _____

CLIENT DEMOGRAPHICS - CONFIRMED: ☐

Name: _____ DOB: _____ (YYYY/MM/DD) Sex: _____

Address: _____

Email: _____ Phone: _____

TREATMENT (as per Public Health Ontario Gonorrhea Testing and Treatment Guide, 2nd Edition)

FIRST-LINE TREATMENT
(Test of cure recommended)

☐ Ceftriaxone 500mg IM Single Dose
Date Prescribed (YYYY/MM/DD): _____

SECOND-LINE TREATMENT (Perform test of cure)

Alternative treatment for anogenital infections

☐ Cefixime 800mg PO + Doxycycline 100mg PO BID for 7 Days
- This regimen is recommended if there is macrolide resistance or contraindication to macrolide use.
- Doxycycline is contraindicated in pregnant and lactating individuals

Date Prescribed (YYYY/MM/DD): _____

Alternative treatment for pharyngeal infections

☐ Cefixime 800mg PO +Azithromycin 1g PO Single Dose

Date Prescribed (YYYY/MM/DD): _____

COMPLETED CLIENT COUNSELLING

☐ Notified of positive result

☐ Advised to abstain from sexual contact for 7 days after treatment

☐ Advised to abstain from sexual contact with untreated sexual contact(s)

☐ Informed that retreatment required if vomiting occurs within 1 hour of taking Azithromycin

☐ Recommended test of cure*

***A test of cure (TOC) is recommended for all positive NG sites in all cases.**
This is particularly important when regimens other than ceftriaxone 500 mg IM are used. Refer to the Test of cure section in the Gonorrhea Guide for more information on the timing for TOC.

- If test of cure is done 3-21 dans after treatment completion, culture is recommended regardless of the patient's symptom status
- If test of cure is done after 21 days from treatment completion and the patient is
  - Asymptomatic-NAAT are recommended
  - Symptomatic-Both culture and NAAT are recommended

CONTACT TRACING (from past 3 months or last know if greater than 3 months)

☐ Client advised to notify sexual contact(s) of need for STI testing and treatment; **AND**

☐ Client advised EOHU will contact them to collect sexual contact(s) information and complete contact tracing

RISK FACTORS

☐ Condom breakage
☐ Diagnosed at time of IUD insertion
☐ Diagnosed at time of therapeutic abortion
☐ More than 1 sexual contact in last 6 months
☐ New sexual contact in past 2 months
☐ No condom used
☐ Opposite sex partner(s)
☐ Parent of a neonate
☐ Pregnant (Number of weeks: _____)
☐ **On PrEP**
☐ Same sex partner(s)
☐ Sex trade worker
☐ Sexually assaulted
☐ Other: _____

REASON FOR TESTING

☐ Contact tracing (sexual contact tested positive)
☐ Symptoms: _____
☐ Routine

NOTES

_____

_____

THE EOHU WILL CONTACT ALL CLIENTS INFECTED WITH GONORRHEA.

13126 ICD-26e

If you require this information in an alternate format, please call 1-800-267-7120 and press 0.